

Emergency Plan

The following are signs/symptoms of an allergic reaction:

- Mouth – Itching, tingling, or swelling of lips, tongue, mouth
- Skin – Hives, itchy rash, swelling of the face or extremities
- GI – Nausea, abdominal cramps, vomiting, diarrhea
- Throat – Tightening of throat, hoarseness, hacking cough
- Lungs—Shortness of breath, repetitive coughing, wheezing
- Heart -- Weak or thready pulse, fainting, pale, blueness

The severity of symptoms can quickly change. All above symptoms can potentially progress to a life-threatening reaction.

1. If student experiences a sting for which he/she has history of severe reaction, do the following: (Check all those that apply and indicate any medication that should be given)

- Attempt to scrape off stinger
- Give the following medication(s), as indicated:

(Name of Medication)	(Amount/Dosage)	(Indication/Timing [immediately or when specific symptoms occur or progress])
(Name of Medication)	(Amount/Dosage)	(Indication/Timing [immediately or when specific symptoms occur or progress])
(Name of Medication)	(Amount/Dosage)	(Indication/Timing [immediately or when specific symptoms occur or progress])

- Apply ice to the site of the sting

Then make the following emergency calls

2. Call 911. State that an allergic reaction has occurred, and additional epinephrine may be needed.

3. Dr. _____ Phone Number _____

4. Parent _____ Phone Number(s) _____

Other Emergency Contacts:

Name/Relationship

Phone Numbers

EVEN IF PARENT/GUARDIAN CANNOT BE REACHED, DO NOT HESITATE TO MEDICATE OR TAKE THE STUDENT TO MEDICAL FACILITY

Parent/Guardian Signature _____ Date _____

Doctor's Signature _____ Date _____

(Note: If an Epi-Pen are injected as treatment for a bee/wasp sting, EMS (911) will be called as per recommended guidelines as additional epinephrine may be needed.)